

Polar amplification of Pliocene climate by elevated trace gas radiative forcing

Hopcroft, Peter; Ramstein, Gilles; Pugh, Thomas; Hunter, Stephen; Murguia-Flores, Fabiola; Quiquet, Aurelien; Sun, Yong; Tan, Ning; Valdes, Paul J.

DOI:
[10.1073/pnas.2002320117](https://doi.org/10.1073/pnas.2002320117)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Hopcroft, P, Ramstein, G, Pugh, T, Hunter, S, Murguia-Flores, F, Quiquet, A, Sun, Y, Tan, N & Valdes, PJ 2020, 'Polar amplification of Pliocene climate by elevated trace gas radiative forcing', *National Academy of Sciences. Proceedings*, vol. 117, no. 38, pp. 23401-23407. <https://doi.org/10.1073/pnas.2002320117>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Polar amplification of Pliocene climate by elevated trace gas radiative forcing

Peter O. Hopcroft^{a,1}, Gilles Ramstein^b, Thomas A.M. Pugh^{a,c}, Stephen J. Hunter^d, Fabiola Murguía-Flores^e, Aurélien Quiquet^{b,f}, Yong Sun^{b,g}, Ning Tan^{b,h}, and Paul J. Valdes^e

^aSchool of Geography, Earth & Environmental Sciences, University of Birmingham, U.K. B15 2TT; ^bLaboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.; ^cBirmingham Institute of Forest Research, University of Birmingham, U.K. B15 2TT; ^dSchool of Earth and Environment, University of Leeds, Leeds, UK.; ^eBristol Research Initiative for the Dynamic Global Environment, School of Geographical Sciences, University of Bristol, U.K. BS8 1SS.; ^fInstitut Louis Bachelier, Chair Energy and Prosperity, Paris, 75002, France.; ^gState Key Laboratory of Numerical Modelling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China.; ^hKey Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China.

This manuscript was compiled on July 16, 2020

Warm periods in Earth's history offer opportunities to understand the dynamics of the Earth System under conditions that are similar to those expected in the near future. The mid-Pliocene warm period (mPWP) from 3.3-3.0 million years ago, is the most recent time when atmospheric CO₂ levels were as high as today. However, climate model simulations of the Pliocene underestimate high latitude warming that has been reconstructed from fossil pollen samples and other geological archives. One possible reason for this is that enhanced non-CO₂ trace gas radiative forcing during the Pliocene, including from methane (CH₄), has not been included in modelling. We use a suite of terrestrial biogeochemistry models forced with mPWP climate model simulations from four different climate models, to produce the first comprehensive reconstruction of the mPWP CH₄ cycle, including uncertainty. We simulate an atmospheric CH₄ mixing ratio of 1000-1200 ppbv, which in combination with estimates of radiative forcing from N₂O and O₃, contributes a non-CO₂ radiative forcing of 0.9 Wm⁻² (range 0.6-1.1), which is 43% (range 36-56%) of the CO₂ radiative forcing used in mPWP climate simulations. This additional forcing would cause a global surface temperature increase of 0.6-1.0 °C, with amplified changes at high-latitudes, improving agreement with geological evidence of mid-Pliocene climate. We conclude that natural trace gas feedbacks are critical for interpreting climate warmth during the Pliocene and potentially many other warm phases of the Cenozoic. These results also imply that using Pliocene CO₂ and temperature reconstructions alone, may lead to overestimates of the "fast" or "Charney" climate sensitivity.

Methane | Pliocene | GCM | Trace gas | Biogeochemistry | Wetland

1. Introduction

The mid-Pliocene warm period around 3.3-3.0 million years ago was the last period in Earth's history when atmospheric CO₂ was comparable to today's level, at approximately 400 ppmv (1-4). The mPWP could therefore provide useful information on the response of the Earth System to greenhouse gas-induced warming, that is relevant to the future evolution of the Earth System under continued anthropogenic greenhouse gas emissions (5-9). According to syntheses of reconstructed surface temperatures from geological archives on land (10) and in the ocean (11, 12), the Earth was globally warmer than the pre-industrial, with a significant polar amplification especially in the Northern Hemisphere. The warmer conditions acted to reduce global ice volume, so that sea-level was around 20 m higher than present (13-15).

The Pliocene Model Intercomparison Project (PlioMIP) is a co-ordinated study of climate model responses to Pliocene

boundary conditions (16, 17) aimed at quantifying the underlying drivers of warmth during this time and to better understand the Earth System response to an atmospheric CO₂ concentration of ~400 ppmv. The results from the first phase (PlioMIP1) showed that coupled general circulation models (GCMs) failed to reproduce high-latitude warming seen in reconstructions (16). Several hypotheses could explain why the model simulations underestimate warming, including the role of orbitally induced climate variability (16), the configuration of ocean gateways and palaeogeography (e.g. 18, 19) and radiative forcing from trace gases other than CO₂ (20). Whilst several studies have addressed the first three possibilities, the contribution of the atmospheric methane, nitrous oxide and ozone (CH₄, N₂O, O₃) have not been considered together.

At present CH₄ is the second most important anthropogenic greenhouse gas after CO₂ (21). Its nearly three-fold concentration increase since CE 1750 is responsible for approximately 25% of the greenhouse gas radiative forcing. We know that variations in CH₄ are huge in intensity and abrupt during the Anthropocene relative to the past several thousand years. Before widespread direct atmospheric monitoring, CH₄ could only be traced through air bubbles trapped in polar ice-cores.

Significance Statement

Warm periods in Earth's history provide the only empirical evidence of how the climate system responds to raised atmospheric carbon dioxide (CO₂) levels. The mid-Pliocene, 3.3-3.0 million years ago, was the last time when CO₂ levels were as high as today. However, climate model simulations of the Pliocene underestimate the warming that has been reconstructed from geological archives. Using a numerical model of the global methane cycle we show that the inclusion of enhanced concentrations of non-CO₂ trace gases, could have been responsible for an additional warming of 0.6-1.0°C, with larger increases over northern landmasses. These findings demonstrate the importance of trace gas climate forcing both for the Pliocene and potentially warm periods during much of Earth's recent history.

POH, GR, and PJV conceived research. POH developed offline models for wetlands, soil NO_x, termites. TP and POH carried out LPJ-GUESS simulations, NT, CC, SJH ran GCM simulations. FM ran soil CH₄ uptake model simulations. All authors contributed to interpreting the results and writing the manuscript.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: p.hopcroft@bham.ac.uk

These ice-core records provide robust evidence that CH₄ is very sensitive to climate, with higher values during warmer periods, and with large-amplitude rapid variations during abrupt climate events (22, 23). Nitrous oxide (N₂O) also displays similar characteristics (24), whilst O₃ and many other important atmospheric constituents (e.g. OH) cannot be reconstructed from ice-core gas samples. Before 800,000 years ago, variations in all trace gases with the exception of CO₂, are essentially unknown, and must therefore be simulated using models (e.g. 25).

Beerling et al. (2009) simulated enhanced wetland emissions during the Pliocene but did not link this with other methane sources or sinks(26). Unger & Yue, 2014(20) (UY14 hereafter) simulated an increase in CH₄ lifetime by 23-31% during the Pliocene (for CH₄=1000 - 2000 ppbv) and an increase in tropospheric O₃ by 21-25% which contributed a radiative forcing of approximately 0.3 Wm⁻². However, this CH₄ lifetime increase is partly caused by the self-feedback (e.g. 27), a function of the prescribed CH₄ concentration in the model. Other than a model study of the Eocene and late Cretaceous super-warm periods (28), there is no comprehensive understanding of how trace gases may have affected pre-Quaternary warm periods like the Pliocene.

We address this with an ensemble of terrestrial biogeochemistry simulations for the Pliocene. In combination with climate fields from several GCM simulations of the Pliocene we develop the first estimate of the Pliocene global CH₄ source, CH₄ sinks and thereby the CH₄ concentration, and CH₄ radiative forcing, including uncertainty. We combine these with estimates of N₂O and O₃ radiative forcing to evaluate for the first time how all of these trace gases may have determined the climate during the Pliocene.

2. Results

A. Emissions of CH₄ and Other Trace Gases. Our approach follows that used for modelling of the Quaternary CH₄ variations (25, 29, 30). We drive the LPJ-GUESS dynamic global vegetation model (31, 32) with climate simulations from four models used in the Pliocene Model Intercomparison Project phases 1 and 2 and simulate the key trace gas emissions at the land surface, including CH₄, emissions from wildfires, non-methane biogenic volatile organic compounds (NBVOCs) from vegetation and soil nitrogen oxides (NO_x), see SI Appendix, Fig. S1. Assuming that the Pliocene is a quasi-equilibrium climate state, we do not account for any increased emissions from marine clathrates or permafrost which are likely more sensitive to abrupt warming (33). We use PlioMIP phase 1 (experiment 2) simulations (34) with CCSM4, GISS-E2-R and IPSL-CM5A-LR and PlioMIP phase 2 (Eoi⁴⁰⁰) (16) simulations with IPSL-CM5A-LR and HadCM3-M21. In all Pliocene climate model simulations the orography, land-ice and vegetation are based on geological reconstructions for the Pliocene (35, 36) and the mixing ratio of atmospheric CO₂ is increased from 280 ppbv in the pre-industrial to 400 or 405 ppbv. We also incorporate HadCM3 PlioMIP phase 2 simulations performed with atmospheric CO₂ mixing ratios of 450 (50) and 490 ppmv which are labelled Eoi⁴⁵⁰ and Eoi⁴⁹⁰, respectively. All climate simulations are described in Materials and Methods.

We estimate the CH₄ lifetime and the resultant concentration of CH₄ using a simplified offline one-box model of the

atmospheric CH₄ chemistry (37) and radiative forcing (38). With this we do not resolve atmospheric transport or detailed chemical pathways, but we are able to quantify uncertainty by sampling the model stochastically. See Methods and Supporting Information SI Appendix, Table S2 for full details of all models used.

All CH₄ emissions increase in response to the simulated Pliocene climate conditions (see SI Appendix, Table S1). Despite warmer and in many places wetter conditions (see SI Appendix, Fig. S2), wetland area increases only marginally, but the emissions increase by 20-46%. Emissions increase notably in the Sahel region of North Africa in all models, and in South Africa in all models except GISS. Elsewhere, emissions increase in South East Asia in all models except GISS and Australia in all models except CCSM4 and GISS. GISS shows the largest emissions increase of all models over the Amazon, whereas both IPSL simulations show a slight reduction in this region. A factor separation approach summarised in SI Appendix, Table S3, demonstrates that increased Pliocene temperature enhance emissions in all models. In HadCM3 and GISS wetter soils also promote emissions, but are less important in the other GCMs. Soil carbon stocks act to reduce emissions in HadCM3, and increase emissions in CCSM4, but have much less impact in the other models. Wildfire, and termite emissions also increase by 50-83% and 21-37%, respectively and soil uptake (at pCH₄=715 ppmv) increases by 8-27%. Isoprene emissions increase by 5-40% and monoterpene changes by -9 to +7%. These ranges span the different climate models used to drive each emissions scheme.

B. CH₄ Chemical Lifetime and Radiative Forcing. Although increased emissions will translate into elevated atmospheric concentration, the concentration is also dependent on the lifetime, which is influenced by other trace gas emissions and the physical state of the atmosphere. The change in CH₄ lifetime, Pliocene minus pre-industrial, is summarised for each GCM in figure 1. Increased CH₄ emissions from wetlands, wildfires and termites contribute a positive lifetime anomaly through the self-feedback effect. Additional NMVOC emissions (CO and isoprene) from wildfires and vegetation enhance the lifetime slightly. The higher global temperatures and associated humidity increase act to offset some of this lifetime increase in each model. This is because the reaction rate of CH₄ with OH scales with temperature and the production of OH which oxidises CH₄, increases with water vapour availability and hence temperature (e.g. 37, 39). BVOCs are the most model-dependent term, with a strong increase in emissions during the Pliocene in IPSL and HadCM3 but not in the other two GCMs. This is most likely a result of the differences in the response of the tropical hydrological cycle. Soil NO_x shows agreement across the four climate models. The results are also in agreement with the results of UY14. Lightning NO_x increase and increased O₃ in the troposphere (both taken from UY14) have opposing influences on the Pliocene lifetime.

As the CH₄ lifetime is dependent on the CH₄ level, there is a non-linearity which means that the total change in lifetime does not equal the sum of these individual terms. The net change is positive for HadCM3 and both IPSL simulations and very weakly negative in the other two models. This is mostly due to the much stronger BVOC emission term in the former two models.

The results of the sampling of the total Pliocene minus

pre-industrial CH₄ radiative forcing are shown in figure 2. The mean \pm 1 standard deviation anomalies are 0.16 \pm 0.02, 0.20 \pm 0.03, 0.21 \pm 0.03 and 0.33 \pm 0.05 Wm⁻² for CCSM4, GISS, IPSL and HadCM3, respectively. The combined mean and 1 σ range is 0.22 \pm 0.07 Wm⁻². The total uncertainty in this estimate is dominated by the choice of climate simulation.

3. Discussion

A. Total Radiative forcing. The global mean radiative forcing due to the CO₂ concentration of 400 ppm during the Pliocene is around 2.0 \pm 0.3 Wm⁻². We approximate N₂O radiative forcing based on natural variations in these three greenhouse gases through the late Quaternary as 15% of the combined CO₂ and CH₄ radiative forcing(40). Radiative forcing by O₃ is 0.29 Wm⁻² as simulated for CH₄=1000 ppbv by UY14(20). This positive radiative forcing is partly due to a simulated increase in emissions of O₃ precursors, consistent with our simulated emissions (SI Appendix, Table S1). The uncertainty on the O₃ radiative forcing is calculated by scaling with the CH₄ concentration, which in three sensitivity simulations by (20) shows an approximately logarithmic dependence. O₃ is also created as a by-product of CH₄ emissions, thus it is partially included in the O₃ radiative forcing and in our indirect CH₄ radiative forcing. Hence we reduce our total forcing using the combined factor for O₃ radiative forcing from CH₄ as in SI Appendix, Table S4, to avoid double counting. Together with a CH₄ radiative forcing, we derive an approximate radiative forcing due to other non-CO₂ well-mixed GHGs of 0.8 (0.62-1.02) Wm⁻² as summarized in table 1.

The rate of methane radiative forcing increase per degree of global mean temperature change (see also 41) is a useful property to compare time periods or models. We denoted this as γ_{fCH_4} and it ranges from 89-113 mWm⁻²/K across the 5 climate model simulations, with the lowest value simulated by CCSM4 and the highest by HadCM3. This range slightly exceeds the observed value of 68 (57-85) mWm⁻²/K derived from the last glacial maximum to the late-Holocene pre-industrial (22, 42). It is consistent with simulated values of 97-119 mWm⁻²/K for the Early Eocene and 89 mWm⁻²/K for the Late Cretaceous calculated with a coupled climate-chemistry modelling framework (28), for which the direct radiative forcing values have been augmented here following a recent update (38).

The Pliocene climate may have induced changes in the emissions and lifetime of natural aerosols. UY14 simulated a net cooling by nitrate, particulate organic matter and biogenic secondary organic aerosols (SOA) and a warming effect from black carbon, leading to a negative aerosol forcing of -0.4 Wm⁻² (20). We simulate smaller changes in emissions from biomass burning (+50-84% versus a +101% by UY14) and BVOCs (precursors of SOA: +5-40% versus +50% by UY14), and so this would likely equate to a smaller net forcing. Other aerosol changes may have a warming effect. For example mineral dust levels are generally lower in warm climates, and deserts contracted during the Pliocene (10). Interactions of CH₄ with nitrate and sulphate can also enhance radiative forcing (43). Pliocene aerosol effects therefore require further study.

The climate simulations could be benchmarked with Pliocene climate reconstructions. Though no global compilations of precipitation change are available for the Pliocene,

individual records can provide informative constraints. The West African monsoon was probably stronger during the mid-Pliocene even when compared to the early to mid-Holocene (44). There was also a general drying trend in East Africa during the Pliocene (45), which if continued to the present, implies wetter conditions during the mid-Pliocene. Both of these features are most faithfully reproduced in HadCM3 and GISS and the Eoi⁴⁰⁰ IPSL simulation (see SI Appendix, Fig. S2). In Asia, a reconstructed near-doubling of precipitation in Southern China (46) is replicated in IPSL (Eoi⁴⁰⁰ simulation) and by HadCM3 but not the other models. Only HadCM3 and CCSM4 simulate wetter conditions reconstructed for the early to mid-Pliocene in Australia (47). One of the regions of largest inter-model spread in precipitation is tropical South America, but to our knowledge quantitative precipitation records have yet to be produced.

HadCM3 has the strongest warming globally and in particular over most land masses and over the North Atlantic, where most other models significantly underestimate reconstructed warming (6). HadCM3 model also shows among the best agreement with the available temperature reconstructions(10). This lends support to the stronger methane forcing that is simulated with the HadCM3 climate drivers, since the enhanced warming, particularly over land and associated hydrological cycle changes, promote trace gas emissions. A global biome reconstruction could also provide some measure to discern between simulations (48), but this is initially derived from a simulation with HadCM3, and so is very likely biased towards the climatic response of this model.

B. Likely non-CO₂ Climatic Response. Our modelling study probably underestimates the true magnitude of the Pliocene methane cycle feedback, because the climate and methane cycle are not coupled. The feedback of methane radiative forcing via climate to methane emissions is not complete, and all of the simulated climates underestimate mid- and high-latitude warming over land considerably (10). Furthermore, our study has a relatively modest change in wetland area compared with modelling studies of earlier warm periods (26, 28, 49), and so the resultant radiative forcing is potentially a conservative prediction. Any additional warming resulting from the higher concentrations of trace gases including methane, would further perturb the sources and atmospheric chemistry. Using the upper and lower limits for the γ_{fCH_4} , we can approximate this. The trace gas forcing-induced warming would lead to an additional methane feedback bringing the total Pliocene methane radiative forcing to 0.62-1.1 Wm⁻². These total non-CO₂ values are 33-56% of the CO₂ radiative forcing, with a central estimate of 0.9 Wm⁻² or 43% of the CO₂ forcing. This would cause a warming of 0.6-1.0 °C (see table 1).

This is an important additional warming signal given that PlioMIP phase 1 GCMs forced only with increased CO₂ show a response of 2.7 \pm 0.8 K (16). To better understand the regional impacts of this additional radiative forcing we analyse mid-Pliocene simulations from HadCM3 with prescribed levels of atmospheric CO₂. The CO₂ radiative forcing differences relative to the default Eoi⁴⁰⁰ are approximately 0.66 Wm⁻² for the increase by 50 ppmv and 1.1 Wm⁻² for the increase to 490 ppmv. These encompass the range of non-CO₂ GHG radiative forcing we calculated (0.62-1.1 Wm²). The effective radiative forcing (or the temperature change per unit increment of radiative forcing) is actually higher for CH₄ and N₂O than

for CO₂, and it is lower for O₃ (51). For simplicity here we assume that they are all equal.

The Pliocene HadCM3 simulations with differing CO₂ concentrations show significant polar amplification, especially in the Northern hemisphere and around the Atlantic see figure 3. The Eoi⁴⁵⁰ and Eoi⁴⁹⁰ simulations show relatively muted warming signal in the circum-Atlantic, because of a reduction in heat transport in the Atlantic (50). The upper-end of our estimated radiative forcing, represented by the Eoi⁴⁹⁰ simulation, is shown in figure 3b. This estimated non-CO₂ GHG radiative forcing (figure 3c) causes high-latitude temperatures to increase by around 1-2.5°C over land and by 1-2°C over the ocean surface (figure 3 and see SI Appendix, Fig. S3). This does not eliminate the large mismatches found over high latitude Asia (figure 3c,d) and in the North Atlantic (see SI Appendix, Fig. S3), but it improves the comparison with the temperature reconstructions by (10) as shown in figure 3d. The distribution of errors is shifted from having a significant probability over the range -5 to -1°C to being approximately centred on 0°C. For HadCM3, both the smaller and larger increases in radiative forcing improve the model-data agreement, except for sites with very high temperatures at high-latitudes. The very high temperatures in high-latitude regions would very likely further enhance trace gas emissions, especially CH₄ from wetlands, and so reinforce the positive radiative forcing.

Reasons for the enduring high-latitude discrepancies might include seasonal bias in reconstructions over land or ocean (52), and long-term trends in seawater chemistry (53). If the reconstructions capture peak warmth during orbital cycles, this may not be adequately captured by climate model simulations with pre-industrial orbital configuration (54), and global assemblages of reconstructions may capture different phases of Earth's orbit in different locations (10), further complicating comparisons with simulations. Terrestrial reconstructions are also potentially influenced by the dependence of plant water-use efficiency on atmospheric CO₂, and changes in seasonality and the frequency of extreme events (55). However, a general underestimation in polar amplification of past warm states, could signify systematic problems with climate model parameterisations of clouds (56, 57) and/or aerosol-cloud interactions (58).

These model results allow an estimate of $S_{LI,GHG}$, the temperature response per unit of radiative forcing due to ice-sheets and sea-level (LI) and due to greenhouse gases (GHG) (K/Wm⁻²), where S_{LI,CO_2} was possible until now (4). Several studies have considered the Earth System Sensitivity, the long-term temperature response to changes in radiative forcing, incorporating vegetation and ice-sheet responses (5). Our central estimate of 43% non-CO₂ greenhouse gas radiative forcing would reduce values estimated from Pliocene temperature reconstructions from around 9°C (59) to 6.5°C. This is still much larger than predicted based on "fast" feedbacks in the climate system, and is therefore potentially consistent with irreversible long-term planetary warming, should we fail to limit warming this century to 1.5°C (8).

4. Conclusions

The mid-Pliocene is a critically important past time period to understand because levels of atmospheric CO₂ were very likely as high as today's anthropogenically perturbed levels, at around 400 ppmv. The Pliocene therefore offers unique

insight into a warmer Earth system at equilibrium (5, 6). CH₄ is the second most important anthropogenic greenhouse gas today, but beyond the ice-core era to 800 ka BP, it has been largely ignored. Whereas for CO₂, proxies have been developed to reconstruct its evolution for deep times, no such proxy exists for CH₄. We used vegetation model simulations and an offline CH₄ budget model to produce a comprehensive set of simulations quantifying the likely CH₄ emissions, lifetime and concentration during this time period, investigating changes of methane sources and sinks but excluding abrupt releases derived from clathrates or permafrost degradation. We show that there is a direct net positive radiative forcing of 0.22 Wm⁻² (range 0.1-0.45), which when combined with estimates of radiative forcing from N₂O and O₃ leads to an additional radiative forcing that is between 36-56% of that caused by a CO₂ concentration of 400 pmv. This would likely cause an additional global mean warming of approximately 0.6-1.0 °C at the global scale. Terrestrial trace gas radiative forcing is therefore critically important for understanding the reconstructed warmth of the Pliocene. We suggest that the first-order estimates of these additional forcing agents could help to reconcile model and proxy-based estimates of Pliocene warmth and that failing to include such forcings will lead to substantial biases in simulations of past climates. Their omission also has consequences when trying to estimate the "fast" or "Charney" climate sensitivity from paleo-data (60). Such methods attempt to remove the slower "Earth System" forcings such as changes in ice sheets and assume the residual changes are due to the fast response to CO₂. However, these estimates have not normally included non-CO₂ trace gases and are therefore potentially overestimating the CO₂ sensitivity.




Fig. 1. Simulated mean change in lifetime of CH₄ with respect to OH for the Pliocene minus the pre-industrial, net values and individual terms are shown for the five different sets of climate drivers (HadCM3-M21-Eoi⁴⁰⁰, IPSL-CM5A-PlioMIP1, ISPL-CM5A-Eoi⁴⁰⁰, GISS-E2-R-PlioMIP1, CCSM4-PlioMIP1).

373 **Figures.**




Fig. 2. Estimated radiative forcing expressed as a probability density function. The uncertainty stems from (i) assumed $\pm 20\%$ uncertainties in lightning NO_x and tropospheric O_3 (ii) uncertainty in the parameters used to estimate OH lifetime of CH_4 (37), and (iii) uncertainty in the total radiative forcing due to CH_4 (37, 38). The $\pm 1 \sigma$ uncertainty is shown above each curve. Dashed lines represent first two factors and solid bars underneath include all three sources of uncertainty.




Fig. 3. Simulations with HadCM3-M21 and reconstructions of Pliocene near-surface air temperature change (ΔT) relative to the pre-industrial (K). (a) Simulated (50) and reconstructed (10) ΔT for Eoi⁴⁹⁰ minus pre-industrial, (b) simulated ΔT for Eoi⁴⁹⁰ - Eoi⁴⁰⁰, (c) latitudinal comparison of the reconstructed and simulated temperature anomalies, (d) histogram of model minus reconstruction temperature anomalies.

Table 1. Pliocene radiative forcing, CH₄ cycle sensitivity and predicted non-CO₂-forced warming from this study and past work.

| | Mean | Range | Reference |
|---|------|-----------|--|
| Radiative forcing (Wm⁻²) | | | |
| CO ₂ | 2.0 | - | Haywood et al. 2013(17) |
| CH ₄ | 0.22 | 0.10-0.43 | This study |
| N ₂ O | 0.33 | 0.32-0.36 | This study |
| O ₃ | 0.29 | 0.28-0.36 | Unger & Yue, 2014(20) |
| Total GHG | 2.80 | 2.67-3.02 | This study |
| Total non-CO ₂ | 0.80 | 0.67-1.02 | This study |
| with self-feedback | 0.86 | 0.62-1.12 | This study |
| % of CO ₂ | 43 % | 36-56 % | This study |
| CH₄ cycle sensitivity to temperature (mWm⁻²K⁻¹) | | | |
| Pliocene (3.3-3.0 Ma BP) | 100 | 89-114 | This study |
| Early Eocene (55 Ma BP) | 109 | 97-120 | Beerling et al. 2011(28) |
| Cretaceous (90 Ma BP) | 89 | - | Beerling et al. 2011(28) |
| LGM (21 ka BP) | 68 | 57-85 | Loulergue et al. 2008; Annan & Hargreaves, 2013 (22, 42) |
| Global mean temperature change (°C) | | | |
| Warming (non-CO ₂) | 0.7 | 0.6-0.9 | This study |
| with self-feedback | 0.7 | 0.6-1.0 | This study |

375 **Materials and Methods**

376

In PlioMIP1 phase 1, the boundary conditions are referred to as experiment 2 for coupled atmosphere-ocean simulations (34, 61). In PlioMIP phase 2 the nomenclature used is Ex^c, where x can include 'o' and/or 'i' to signify whether or not the orography and land-ice respectively are based on Pliocene geological reconstructions, and 'c' is the prescribed Pliocene CO₂ mixing ratio (17). Since the default PlioMIP phase 2 CO₂ mixing ratio is 400 ppmv, this simulation is labelled Eoi⁴⁰⁰. We also make use of simulations with HadCM3-M21 in which the CO₂ mixing ratio is set to 450 or 490 ppmv, and these are labelled Eoi⁴⁵⁰ and Eoi⁴⁹⁰, respectively. The global mean radiative forcing difference relative to the standard Eoi⁴⁰⁰ simulation is therefore approximately 0.7 and 1.1 Wm⁻², for Eoi⁴⁵⁰ and Eoi⁴⁹⁰, respectively. These increases therefore encompass the upper and lower end of our estimated additional non-CO₂ radiative forcing listed in table 1.

A. Climate simulations. We use climate simulations from four GCMs: HadCM3-M21, IPSL-CM5A-LR, CCSM4 and GISS-E2-R. These GCMs are compared briefly in SI Appendix, Table S2. They span a reasonable range of changes for Pliocene and incorporate either PlioMIP phase 1 experiment 2 (34, 61) or PlioMIP phase 2 boundary conditions for the Pliocene (17). In all Pliocene simulations, the CO₂ mixing ratio is increased from 280 ppmv in the pre-industrial to 400 (PlioMIP 2) or 405 ppmv (PlioMIP1). The remaining Pliocene boundary conditions are based on geological evidence for orography, land-ice, vegetation and soils, or are left unchanged at pre-industrial settings. For land-ice, the volume of the Greenland ice-sheet is reduced and the West Antarctic ice-sheet is removed. Eustatic sea-level is 25 m above modern. The global vegetation reconstruction prescribes a northwards extension of forests causing a reduction in the area of tundra, whilst deserts are partly replaced by savannah and forest, particularly in the Sahel (10). In PlioMIP phase 1 simulations orography increases over the Rockies and East Antarctica, but these are changes are smaller in PlioMIP phase 2 for which both the Canadian Archipelago and Bering Strait are closed. The remaining model boundary conditions, including orbital configuration, solar constant, mixing ratios of CH₄, N₂O and O₃ and aerosols are as used in the relevant pre-industrial setup.

B. Vegetation model simulations. We simulate the land surface with the dynamic stand-based global vegetation model LPJ-GUESS v3 (31, 32). LPJ-GUESS was forced with repeating 30 years of monthly fields from the four GCMs listed above. A 500 year spin up to equilibrium was conducting using repeating the 30-year climate data prior to each 30 year simulation. We expand the land-sea-mask using coastlines from PlioMIP phase 2 (17), including a partially deglaciated Greenland. The soil type over new land points is extrapolated from nearby existing points using the default soil type in LPJ-GUESS. CO₂ was set to 280 and 400 ppmv for pre-industrial and Pliocene simulations, respectively.

C. Trace gas emissions. Wetland area is calculated from the GCM soil moisture using a TOPMODEL approach (62), in which the fractional area of inundation is calculated from the probability density function within each gridcell as derived from global high-resolution topographic data. CH₄ emissions are a function of microbial activity i.e. temperature, available substrate and wetland area (63). Total pre-industrial emissions are scaled to 140 TgCH₄/yr for each model, consistent with the observed methane isotope and concentration measurements (64).

Termite CH₄ emissions are calculated from the LPJ-GUESS simulated plant functional types (PFTs) coverage and observed emission per biome type (30, 65). Isoprene and monoterpene emissions and biomass burning are represented with process-based schemes coupled within LPJ-GUESS (66-68). We estimate soil emissions of nitrogen oxides (NO_x) using a recently developed semi-empirical scheme(69).

D. CH₄ chemical lifetime and radiative forcing. We calculate the uptake of CH₄ in soils with a process-based model (70) driven with climatologies from the GCMs. For the OH lifetime τ , we integrate the Pliocene changes in emissions of all of the above species as well as atmospheric conditions and composition. As a full 3D chemistry-transport simulation is extremely computationally expensive, it would limit the evaluation of uncertainty. Hence we employ a parametric model (37) which is calibrated with three coupled climate-chemistry transport models. This model is described in the SI.

We combine the CH₄ source estimates with the lifetime calculation in a global-mean budget calculation $B = S \times \tau$, where B is the atmospheric CH₄ burden in Tg, S is the global mean CH₄ source and τ is the lifetime in years (64). The budget equation is combined with the parametric lifetime model and the soil uptake, which is multiplied by the resultant concentration divided by pre-industrial value. This is solved iteratively, to account for the self-feedback of CH₄ on lifetime and soil uptake.

The methane radiative forcing is a combination of direct and indirect components which are given in the SI Appendix, Table S4. The total is $0.8 \pm 0.09 \text{ Wm}^{-2}$ per 1000 ppbv increase in atmospheric CH₄.

To account for uncertainties, we prescribe a standard deviation of $\pm 20\%$ for lightning NO_x and stratospheric O₃, which are not simulated directly. We include uncertainties inherent in the CH₄ lifetime and radiative forcing calculations as described in the SI and example is shown in SI Appendix, Table S5. The calculations are sampled with 10,000 evaluations of the iterated global CH₄ budget formulation.

E. Data availability. GCM simulation output for the pre-industrial simulations are available from Earth System Grid Federation CMIP5 archive. Pliocene simulations can be obtained through the PlioMIP project. Pliocene geographic boundary conditions used in this study are available from PRISM https://geology.er.usgs.gov/egpsc/prism/4_data.html. LPJ-GUESS model outputs and all model code developed in this study have been archived on figshare: [dx.doi.org/10.6084/m9.figshare.12302201](https://doi.org/10.6084/m9.figshare.12302201) (piControl simulations), [dx.doi.org/10.6084/m9.figshare.12302216](https://doi.org/10.6084/m9.figshare.12302216) (PlioMIP1 simulations), [dx.doi.org/10.6084/m9.figshare.12302228](https://doi.org/10.6084/m9.figshare.12302228) (PlioMIP2 simulations) and [dx.doi.org/10.6084/m9.figshare.12344027](https://doi.org/10.6084/m9.figshare.12344027) (emissions and lifetime code). The source code for LPJ-GUESS v4.0 can be obtained on request through Lund University (web.nateko.lu.se/lpj-guess). HadCM3-M21 Eoi⁴⁹⁰ climate fields are archived on figshare: [dx.doi.org/10.6084/m9.figshare.12630356](https://doi.org/10.6084/m9.figshare.12630356).

ACKNOWLEDGMENTS. POH is supported by a University of Birmingham Fellowship. GR is supported by the French project LEFE "ComPreNdrE" (A2016-992936), French State Program Investissements d'Avenir (managed by ANR), ANR HADOC project, grant ANR-17-CE31-0010 of the French National Research Agency. TAMP acknowledges support from the European Research Council under the European Union Horizon 2020 programme (Grant 758873, TreeMort).

This work was carried out using the computational facilities of the Birmingham Environment for Academic Research - <http://www.bear.bham.ac.uk>. We thank Linda Sohl and Mark Chandler from NASA GISS for providing model outputs and Nan Rosenbloom (NCAR) who performed the original CCSM4 simulations. We thank Sophie Szopa for informative discussions on these topics. This is paper number <x> of the Birmingham Institute of Forest Research.

F. References.

1. Seki O, Foster G, Schmidt D, et al. (2010) Alkenone and boron-based Pliocene pCO₂ records. *Earth Planet Sci Lett* 292:201–211.
2. Bartoli G, Honisch B, Zeebe R (2011) Atmospheric CO₂ decline during the Pliocene intensification of Northern Hemisphere glaciations. *Paleoceanography* 26:PA4213.
3. Badger M, Schmidt D, Mackensen A, Pancost R (2013) High resolution alkenone palaeobarometry indicates relatively stable pCO₂ during the Pliocene (3.3 to 2.8 Ma). *Phil T.R. Soc. A* 371(20130094):371.
4. Martinez-Boti M, et al. (2015) Plio-Pleistocene climate sensitivity evaluated using high-resolution CO₂ records. *Nature* 518:49–54.
5. Lunt D, et al. (2010) Earth system sensitivity inferred from Pliocene modelling and data. *Nature Geosci* 3:60–64.
6. Haywood A, Dowsett H, Dolan A (2016) Integrating geological archives and climate models for the mid-Pliocene warm period. *Nature Comm* 7:10646.
7. Burke K, Williams J, Chandler M, Haywood A, et al. (2018) Pliocene and Eocene provide best analogs for near-future climates. *PNAS* 115(52):13288–13293.
8. Steffen W, Rockstrom J, Richardson K, et al. (2018) Trajectories of the Earth System in the Anthropocene. *PNAS* 115(33):8252–8259.
9. Tan N, et al. (2018) Modeling Greenland ice sheet evolution during the Plio-Pleistocene transition: new constraints for pCO₂ pathway. *Nature Comm* 9:4755.
10. Salzmann U, Dolan A, Haywood A, et al. (2013) Challenges in quantifying Pliocene terrestrial warming revealed by data-model discord. *Nature Climate Change* 3:969–974.
11. Dowsett H, et al. (2013) Sea surface temperature of the mid-Piacenzian ocean: a datamodel comparison. *Scientific Reports* 3.
12. Foley K, Dowsett H (2019) Community sourced mid-Piacenzian sea surface temperature (SST) data. *U.S. Geological Survey data release*, doi: 10.5066/P9YP3DTV.
13. Miller K, et al. (2012) High tide of the warm Pliocene: implications of global sea level for Antarctic deglaciation. *Geology* 40:407–410.
14. Dutton A, Carlson A, Lnog A, Milne G, et al. (2015) Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science* 349(6244):aaa4019–1.
15. Dumitru O, Austermann J, Polyak V, Fornos J, et al. (2019) Constraints on global mean sea level during Pliocene warmth. *Nature* 574:233–236.
16. Haywood A, et al. (2013) Large-scale features of Pliocene climate: results from the Pliocene Model Intercomparison Project. *Clim Past* 9:191–209.
17. Haywood A, Dowsett H, Dolan A, Rowley D, et al. (2016) The Pliocene Model Intercomparison Project (PlioMIP) Phase 2: scientific objectives and experimental design. *Clim Past* 12:663–675.
18. Robinson M, et al. (2011) Bathymetric controls on Pliocene North Atlantic and Arctic sea surface temperature and deepwater production. *Palaeogeog, Palaeoclim, Palaeoecol* 309(1–2):92–97.
19. Otto-Bliesner B, Braconnot P, Harrison S, Lunt D, et al. (2017) Two Interglacials: Scientific Objectives and Experimental Designs for CMIP6 and PMIP4 Holocene and Last Interglacial Simulations. *Geosci Model Dev* p. .
20. Unger N, Yue X (2014) Strong chemistry-climate feedbacks in the Pliocene. *Geophys Res Lett* 41(527–533).
21. Myhre G, et al. (2013) Anthropogenic and Natural Radiative Forcing in *Climate Change 2013: The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker T, et al. (C.U.P., Cambridge, U.K. and New York, U.S.A.), pp. 659–740.
22. Loulergue L, et al. (2008) Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years. *Nature* 453(7383):383–386.
23. Rhodes R, et al. (2015) Enhanced tropical methane production in response to iceberg discharge in the North Atlantic. *Science* 348:1016–1019.
24. Schilt A, et al. (2010) Glacial-interglacial and millennial scale variations in the atmospheric nitrous oxide concentration during the last 800,000 years. *Quaternary Sci. Rev.* 29:182–192.
25. Valdes P, Beerling D, Johnson C (2005) The ice age methane budget. *Geophys. Res. Lett.* 32(L02704).
26. Beerling D, Berner R, Mackenzie F, Harfoot M, Pyle J (2009) Methane and the CH₄-related greenhouse effect over the past 400 millions years. *American Journal of Science* 309:97–113.
27. Holmes C (2018) Methane Feedback on Atmospheric Chemistry: Methods, Models, and Mechanisms. *J Advances in Modeling Earth Systems* 10:1087–1099.
28. Beerling D, Fox A, Stevenson D, Valdes P (2011) Enhanced chemistry-climate feedbacks in past greenhouse worlds. *Proc Natl Acad Sci USA* 108:9880–9775.
29. Singarayer J, Valdes P, Friedlingstein P, Nelson S, Beerling D (2011) Late Holocene methane rise caused by orbitally controlled increase in tropical sources. *Nature* 470:82–85.
30. Hopcroft P, Valdes P, O'Connor F, Kaplan J, Beerling D (2017) Understanding the glacial methane cycle. *Nature Communications* 8(14383).
31. Smith B, Prentice I, Sykes M (2001) Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space. *Global Ecology and Biogeog* 10:621–637.
32. Smith B, et al. (2014) Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences* 11:2027–2054.
33. Ruppel C, Kessler J (2017) The interaction of climate change and methane hydrates. *Rev Geophysics* 55:126–168.
34. Haywood A, Dowsett H, Otto-Bliesner B, et al. (2011) Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 2). *Geosci Model Dev* 4:571–577.
35. Dowsett H, Robinson M, Haywood A, et al. (2010) The PRISM3D paleoenvironmental reconstruction. *Stratigraphy* 7:123–139.
36. Dowsett H, Dolan A, Rowley D, et al. (2016) The PRISM4 (mid-Piacenzian) paleoenvironmental reconstruction. *Clim Past* 12:1519–1538.
37. Holmes C, Prather M, Sovde O, Myhre G (2013) Future methane, hydroxyl, and their uncertainties: key climate and emission parameters for future predictions. *Atmos Chem Phys* 13:285–302.
38. Etminan M, Myhre G, Highwood E, Shine K (2017) Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision of the methane radiative forcing. *Geophys Res Lett* 43:12614–12623.
39. Voulgarakis A, et al. (2013) Analysis of present day and future OH and methane lifetime in the ACCMIP simulations. *Atmos Chem Phys* 13:2563–2587.
40. Hansen J, Sato M, Kharecha P, et al. (2007) Climate change and trace gases. *Phil Trans R Soc A* 365(1856):1925–1954.
41. Ringeval B, Friedlingstein P, Koven C, et al. (2011) Climate-CH₄ feedback from wetlands and its interaction with the climate-CO₂ feedback. *Biogeosciences* 8:2137–2157.
42. Annan J, Hargreaves J (2013) A new global reconstruction of temperature changes at the Last Glacial Maximum. *Climate of the Past* 9:367–376.
43. Shindell D, et al. (2009) Improved attribution of climate forcing to emissions. *Science* 326:716–718.
44. Kuechler R, Dupont L, Schefuss E (2018) Hybrid insolation forcing of Pliocene monsoon dynamics in West Africa. *Clim Past* 14:73–84.
45. Liddy H, Feakins S, Tierney J (2016) Cooling and drying in northeast Africa across the Pliocene. *Earth Planetary Science Letters* 449:430–438.
46. Su T, Jacques F, Spicer R, et al. (2013) Post-Pliocene establishment of the present monsoonal climate in SW China: evidence from the late Pliocene Longmen megafloora. *Clim Past* 9:1911–19020.
47. Christensen B, Renema W, Henderiks J, et al. (2017) Cooling and drying in northeast Africa across the Pliocene. *Geophys Res Lett* 44:6914–6925.
48. Salzmann U, Haywood A, Lunt D, Valdes P, Hill D (2008) A new global biome reconstruction and data-model comparison for the Middle Pliocene. *Global Ecol Biogeogr* 17:432–447.
49. Wilton D, Badger M, Kantzas E, et al. (2019) A predictive algorithm for wetlands in deep time paleoclimate models. *Geosci Model Dev* 12:1351–1364.
50. Hunter S, Haywood A, Dolan A, Tindall J (2019) The HadCM3 contribution to PlioMIP Phase 2. *Clim Past* 15:1691–1713.
51. Hansen J (2005) Efficacy of climate forcings. *J Geophys Res* 110:D18104.
52. Tierney J, Tingley M (2016) BAYSPLINE: A New Calibration for the Alkenone Paleothermometer. *Paleoceanography and Paleoclimatology* 33:281–301.
53. Evans D, Brierley C, Raymo M, et al. (2016) Planktic foraminifera shell chemistry response to seawater chemistry: Pliocene–Pleistocene seawater Mg/Ca, temperature and sea level change. *Earth Planet Sci Lett* 438:139–148.
54. Prescott C, Haywood A, Dolan A, et al. (2014) Assessing orbitally-forced interglacial climate variability during the mid-Pliocene Warm Period. *Earth Planet Sci Lett* 400:261–271.
55. Harrison S, Bartlein P, Prentice I (2016) What have we learnt from palaeoclimate simulations? *J Quat Sci* 31(4):364–385.
56. Sagoo N, Valdes P, Flecker R, Gregoire L (2013) The Early Eocene equable climate problem: can perturbations of climate model parameters identify possible solutions? *Phil Trans Roy Soc A* 371(2001):20130123.
57. Zhu J, Poulsen C, Tierney J (2019) Simulation of Eocene extreme warmth and high climate sensitivity through cloud feedbacks. *Science Advances* 5(9):eaax1874.
58. Kiehl J, Shields C (2013) Sensitivity of the palaeocen-eocene thermal maximum climate to cloud properties. *Phil Trans R Soc A* 371.
59. Pagani M, Liu Z, LaRiviere J, Ravel A (2010) High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations. *Nature Geoscience* 3:27–30.
60. PALEOSENS Project Members (2012) Making sense of palaeoclimate sensitivity. *Nature* 491:683–691.
61. Haywood A, Dowsett H, Otto-Bliesner B, et al. (2010) Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 1). *Geosci Model Dev* 3:227–242.
62. Zhang Z, Zimmermann N, Kaplan J, Poulter B (2016) Modeling spatiotemporal dynamics of global wetlands: comprehensive evaluation of a new sub-grid TOPMODEL parameterization and uncertainties. *Biogeosciences* 13:1387–1408.
63. Gedney N, Cox P, Huntingford C (2004) Climate feedback from wetland methane emissions. *Geophys. Res. Lett.* 31(L20503).
64. Hopcroft P, Valdes P, Kaplan J (2018) Bayesian analysis of the glacial-interglacial methane increase constrained by stable isotopes and Earth System modelling. *Geophys Res Lett* 45:3653–3663.
65. Sanderson M (1996) Biomass of termites and their emission of methane and carbon dioxide: A global database. *Glob Biogeochem Cy* 10(4):543–557.
66. Arneeth A, et al. (2007) CO₂ inhibition of global terrestrial isoprene emissions: Potential implications for atmospheric chemistry. *Geophys. Res. Lett.* 34(L18813).
67. Schurgers G, Arneeth A, Holzinger R, Goldstein A (2008) Process-based modelling of biogenic monoterpene emissions: Sensitivity to temperature and light. *Atmos Chem Phys* 8:243–257.
68. Thonicke K, Venevsky S, Sitch S, Cramer W (2001) The role of fire disturbance for global vegetation dynamics: coupling fire into a Dynamic Global Vegetation Model. *Global Ecology and Biogeog* 10:661–677.
69. Hudman R, et al. (2012) Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints. *Atmos Chem Phys* 12:7779–7795.
70. Murguia-Flores F, Arndt S, Ganesan A, et al. (2018) Soil Methanotrophy Model (MeMo v1.0): a process-based model to quantify global uptake of atmospheric methane by soil. *Geosci Model Dev* 11:2009–2032.